

# MCCQ

①

\* Anxiety signs may occur in the following

- except -
- hypoglycemia
  - pheochromocytoma
  - hyperthyroidism
  - Conn's syndrome
  - MI / pericarditis

\* Psychiatric exam includes

- Mental state
- physical
- social
- history
- All of above (✓)

\* Delusions can be best defined as:

- false fixed belief that needs specific (✓)
- percept misinterpretation

Representative

- a view point able to be changed when
- convincing evidence

\* Depression occurs in exapt

- Parkinson's DS
- panic  $\downarrow$
- Hypothyroidism
- MV prolapse (✓)

[NB] All GIT  $\downarrow$  say present in depression

ac All types of obsessive exapt.

- Persecution
- Grandiosity
- Infidelity
- ~~Obsessive~~ (✓)

\* Mania is exapt

- Euphoria
- Nihilistic delusions (✓)
- Hyperactivity
- Delusions of grandiosity

\* Schizophrenia is CCC by all except

- Delusional
- Visual hallucinations
- impaired insight
- memory (✓)
- (+ve) symptoms

\* Best tx of ADHD is

- Amphetamine (✓) ↑ DA in the CC that  
↑ the "focus"
- MAOIs
- ECT
- Benzodiazepines

\* Antidepressants include all except

- MAOIs
- SSRIs
- Sertraline (✓) "it's not HTN"
- TCAs

Extrapyramidal side effects of antipsychotics

- Rigidity & tremor (✓)
- Lethargy
- Akinesia
- Dementia

Risk factors of delirium include

- Dehydration
- ICU
- delirium (✓)
- Pain

(NB) Delirium mostly in both extremes of age & not natural

\* Tendency to ↑ the dose of psychoactive substances to get the same effects as before

- Habituation
- Antitoxin
- Tolerance (✓)
- Abuse
- Synergism



the following substances → dependence

except

- Caffeine (✓)
- Cocaine
- Phencyclidine
- Benzodiazepines
- Ephedrine

\* Social phobia associated by

- fear of social  
  - talking in public (✓)  
  - enclosed spaces

\* Pharmacological treatment of Bipolar:

- Carbamazepine
- No information
- Lithium
- All (✓)

$\sqrt{RT}$  is not a function of  $\sqrt{RT}$ 

No cause of extinction except

- Planch. - Dillenium  
 Social Stabia (H)  
 Parnassia saligna  
 Catelania (H) - Parnassia saligna  
 (H) - Parnassia saligna

\* B2D Earthen & cobblestone unit  
active though

a Diff. bet observed & computed

- Acts, which the pts have to do to alleviate their anxiety (an)

Adams

\* ECT used in all except

Primary disorders (✓)

\* Pharmacotherapy of dep:

- BZD
- SSRI (✓)
- A/B

\* use of Lithium

- Phenothiazine (✓)
- Anticholinergic
- BZD

\* Suicide is more common in except

- post partum psychosis
- severe depression
- previous suicidal attempts
- comorbid disorder (✓)

\* Antipsychotic act:

- Blocking dopamine (✓)

\* Depressive is acc by except

- ↓ self confidence (✓)
- Guilt feeling
- lack of interest
- suicidal

\* Acute dystonia is anti cholinergic  
can be the key:

- A.T. hist
- A.T. chol
- BZD
- All (✓) if emergency
- None

\* Post traumatic stress disorder except

- the person is major & active  
normal human experience

Flash back

- No marked occup. impairment (✓)
- Avoidance symptoms



\* The following can produce psychiatric syms.

- CS
- BB
- INF
- CCP
- All (✓)

\* Symp of GAD:

- Apprehension
- tension
- tizzy
- most of time
- All (✓)

\* Schizo is CC by all except

Delusions

- Prod hallucination
- Disorientate to T, P, P (✓)
- formal thought disorder

## Problem Solving

① IBT → out of  $\phi$   
→ Or shopping → needs Hall pass

Ans  $\Rightarrow$  Hypochondriasis

② cat of the blue → fine - some of Paul  
 $\Rightarrow$  Panic disorder

③ urinary retention by  
Cisapride  
 $\Rightarrow$  TCA

④ 57, 55 yrs, extreme easy fatigue  
Lack of interest in everyday activities  
Difficult to sleep  
 $\Rightarrow$  Thyroid.

5. Bizarre behavior  $\Rightarrow$  schizophrenia

6. ♀ alone in room, clothes torn

No response to tongue biting

No angry reaction

$\Rightarrow$  Psychomotor disturbance

7. Aggression directed to 1, 2, 3

$\Rightarrow$  Delirium

prodromal phase of schizophrenia  
"prodromal phase"

acute